

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

*Reports of States and yearly and monthly reports of cities.*

CALIFORNIA—*Los Angeles.*—Month of August, 1897. Estimated population, 103,000. Total deaths, 110, including enteric fever, 3, and phthisis pulmonalis, 19.

*Sacramento.*—Month of August, 1897. Estimated population, 30,000. Total deaths, 26, including phthisis pulmonalis, 1.

*San Francisco (city and county).*—Month of August, 1897. Estimated population, 360,000. Total deaths, 422, including diphtheria, 10; enteric fever, 4; scarlet fever, 1, and phthisis pulmonalis, 50.

CONNECTICUT.—Month of August, 1897. Reports to the State board of health from 168 towns, having an aggregate population of 883,192, show a total of 1,237 deaths, including enteric fever, 15; diphtheria, 14; measles 2; scarlet fever, 5; phthisis pulmonalis, 94, and whooping cough, 12.

*Westport.*—Month of August, 1897. Estimated population, 4,000. Total deaths, 7.

LOUISIANA—*Shreveport.*—Month of August, 1897. Estimated population, 15,000. Total deaths, 44, including phthisis pulmonalis, 5.

MAINE—*Portland.*—Four weeks ended September 4, 1897. Population, 41,500. Total deaths, 67, including diphtheria, 1; measles, 2; whooping cough, 1, and phthisis pulmonalis, 4.

MASSACHUSETTS—*North Attleboro.*—Month of July, 1897. Estimated population, 6,727. Total deaths, 5, including phthisis pulmonalis, 1.

Month of August, 1897. Total deaths, 6.

MICHIGAN.—Reports to the State board of health, Lansing, from 56 observers, for the week ended September 11, 1897, indicate that dysentery and cholera morbus increased and that intermittent fever and bronchitis decreased in area of prevalence. Phthisis pulmonalis was reported present at 187, enteric fever at 46, diphtheria at 22, measles at 14, and whooping cough at 9 places.

MINNESOTA—*Minneapolis.*—Month of August, 1897. Estimated population, 225,602. Total deaths, 164, including diphtheria, 3; whooping cough, 4, and phthisis pulmonalis, 25.

MISSOURI—*St. Louis.*—Month of August, 1897. Estimated population, 600,000—white, 565,000; colored, 35,000. Total deaths, 766, including diphtheria, 10; enteric fever, 18; scarlet fever, 1; whooping cough, 8, and phthisis pulmonalis, 67.

NEW YORK—*Buffalo.*—Month of August, 1897. Estimated population, 360,000. Total deaths, 455, including diphtheria, 8; enteric fever, 3; measles, 1; scarlet fever, 1; whooping cough, 8, and phthisis pulmonalis, 35.

*Yonkers.*—Month of July, 1897. Estimated population, 40,000. Total deaths, 80, including diphtheria, 4, and phthisis pulmonalis, 4.

Month of August, 1897. Estimated population, 40,000. Total deaths, 83, including enteric fever, 5, and phthisis pulmonalis, 3.

OHIO—*Cleveland.*—Month of August, 1897. Estimated population, 350,000. Total deaths, 477, including diphtheria, 4; enteric fever, 12; whooping cough, 4, and phthisis pulmonalis, 12.

*Columbus.*—Month of August, 1897. Estimated population, 135,000. Total deaths, 104, including diphtheria, 1; enteric fever, 4, and phthisis pulmonalis, 20.

OREGON—*Portland.*—Month of July, 1897. Estimated population, 89,000. Total deaths, 79, including whooping cough, 1, and phthisis pulmonalis, 7.

PENNSYLVANIA—*Williamsport.*—Month of August, 1897. Estimated population, 35,000. Total deaths, 21.

VIRGINIA—*Norfolk.*—Month of August, 1897. Estimated population, 45,000—white, 25,710; colored, 19,290. Total deaths, 26, including enteric fever, 2.

*Richmond.*—Month of August, 1897. Estimated population, 103,560—white, 56,440; colored, 47,120. Total deaths, 129—white, 56; colored, 73, including diphtheria, 1; enteric fever, 3, and phthisis pulmonalis, 10.

WASHINGTON—*Seattle.*—Month of August, 1897. Population, 60,000. Total deaths, 37, including diphtheria, 1, and phthisis pulmonalis, 6.

WISCONSIN—*Milwaukee.*—Month of August, 1897. Estimated population, 275,000. Total deaths, 342, including diphtheria, 3; enteric fever, 2, and phthisis pulmonalis, 22.

*Superior.*—Month of August, 1897. Estimated population, 40,549. Total deaths, 30, including enteric fever, 2.